Vaios/Vaios Total Stemmed Shoulder Investigation

Note: This analysis compares the Vaios/Vaios humeral stem/glenoid combination with all other total stemmed shoulder prostheses.

This combination has been identified as having a significantly higher rate of revision. For a detailed explanation of the process used by the Registry that results in identification of prostheses that have a higher than anticipated rate of revision please refer to the Prostheses with Higher than Anticipated Rates of Revision chapter of the most recent AOANJRR Annual Report, https://aoanjrr.sahmri.com/annual-reports-2022.

Note: Procedures using prostheses with no recorded use in 2021 are excluded from the comparator.

TABLE 1

Revision Rate of Primary Total Stemmed Shoulder Replacement

The revision rate of the Vaios/Vaios total stemmed shoulder combination is compared to all other total stemmed shoulder prostheses.

Table 1: Revision Rates of Primary Total Stemmed Shoulder Replacement

| Component | N Revised | N Total | Obs. Years | Revisions/100 Obs. Yrs (95% CI) |
|------------------------------|-----------|---------|------------|------------------------------------|
| Vaios/Vaios | 19 | 36 | 204 | 9.32 (5.61, 14.56) |
| Other Total Stemmed Shoulder | 698 | 8757 | 49188 | 1.42 (1.32, 1.53) |
| TOTAL | 717 | 8793 | 49392 | 1.45 (1.35, 1.56) |

Yearly Cumulative Percent Revision of Primary Total Stemmed Shoulder Replacement

The yearly cumulative percent revision of the Vaios/Vaios total stemmed shoulder combination is compared to all other total stemmed shoulder prostheses.

Table 2: Yearly Cumulative Percent Revision of Primary Total Stemmed Shoulder Replacement

| CDD | 4.1/ | 2.1/ | 2.1/ | 4.V | E.V. |
|------------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| CPR | 1 Yr | 2 Yrs | 3 Yrs | 4 Yrs | 5 Yrs |
| Vaios/Vaios | 13.9 (6.0, 30.2) | 16.7 (7.9, 33.4) | 27.8 (16.0, 45.5) | 33.3 (20.5, 51.2) | 39.1 (25.3, 57.0) |
| Other Total Stemmed Shoulder | 2.9 (2.5, 3.2) | 4.8 (4.3, 5.2) | 5.8 (5.3, 6.4) | 6.5 (6.0, 7.1) | 7.2 (6.6, 7.8) |
| | | | | | |
| CPR | 6 Yrs | 7 Yrs | 8 Yrs | 9 Yrs | 10 Yrs |
| Vaios/Vaios | 45.5 (30.8, 63.4) | 48.7 (33.6, 66.4) | 48.7 (33.6, 66.4) | 57.7 (40.9, 75.5) | 57.7 (40.9, 75.5) |
| Other Total Stemmed Shoulder | 7.8 (7.2, 8.5) | 8.5 (7.9, 9.3) | 9.6 (8.8, 10.4) | 10.5 (9.7, 11.4) | 11.5 (10.6, 12.5) |
| | | | | | |
| CPR | 11 Yrs | 12 Yrs | 13 Yrs | 14 Yrs | 15 Yrs |
| Vaios/Vaios | | | | | |
| Other Total Stemmed Shoulder | 12.6 (11.5, 13.8) | 13.8 (12.6, 15.2) | 15.0 (13.6, 16.6) | 15.4 (13.9, 17.1) | |

FIGURE 1

Vaios/Vaios


Other Total Stemmed Shoulder

Yearly Cumulative Percent Revision of Primary Total Stemmed Shoulder Replacement

The yearly cumulative percent revision of the Vaios/Vaios total stemmed shoulder combination is compared to all other total stemmed shoulder prostheses. In addition, hazard ratios are reported.

Hazard ratios are reported for specific time periods during which the hazard ratio is constant. This is done to enable more specific and valid comparisons of the risk of revision over time. The pattern of variation in risk has important implications with respect to the underlying reasons for any difference.

Figure 1: Cumulative Percent Revision of Primary Total Stemmed Shoulder Replacement



5

1267

0

982

0

742

0

460

0

193

0

37

Note: Prostheses no longer used in 2021 are excluded from the comparator.

12

2145

8

1649

Primary Diagnosis for Revised Primary Total Stemmed Shoulder Replacement

This table identifies the diagnosis of the primary procedure which was subsequently revised. This information is provided as there is a variation on outcome depending on the primary diagnosis. It is therefore important when considering the reasons for a higher than anticipated rate of revision that there is identification of the primary diagnosis. This information should be compared to the primary diagnosis for the revisions of all other total stemmed shoulder prostheses.

Table 3: Primary Diagnosis for Revised Primary Total Stemmed Shoulder Replacement

| | Vaios/Vaios | | Other Total Ster | nmed Shoulder |
|------------------------------|-------------|---------|------------------|---------------|
| Primary Diagnosis | Number | Percent | Number | Percent |
| Osteoarthritis | 18 | 94.7 | 646 | 92.6 |
| Osteonecrosis | | | 14 | 2.0 |
| Fracture | | | 11 | 1.6 |
| Rheumatoid Arthritis | | | 9 | 1.3 |
| Rotator Cuff Arthropathy | | | 9 | 1.3 |
| Other Inflammatory Arthritis | | | 5 | 0.7 |
| Instability | 1 | 5.3 | 4 | 0.6 |
| TOTAL | 19 | 100.0 | 698 | 100.0 |

Reasons for Revision

This is reported in two ways: a percentage of primary procedures revised and as a percentage of all revision procedures.

% Primaries Revised: This shows the proportional contribution of each revision diagnosis as a percentage of the total number of primary procedures. This percentage can be used to approximate the risk of being revised for that diagnosis. Differing percentages between groups, with the same distribution of follow up time, may identify problems of concern.

% Revisions: The number of revisions for each diagnosis is expressed as a percentage of the total number of revisions. This shows the distribution of reasons for revision within a group but cannot be used as a comparison between groups.

Table 4: Primary Total Stemmed Shoulder Replacement - Reason for Revision (Follow-up Limited to 10.6 Years)

| | | Vaios/Vaios | | Other | Other Total Stemmed Shoulder | |
|------------------------------------|--------|------------------------|-------------|--------|------------------------------|-------------|
| Revision Diagnosis | Number | % Primaries Revised | % Revisions | Number | % Primaries Revised | % Revisions |
| Rotator Cuff Insufficiency | 5 | 13.9 | 26.3 | 229 | 2.6 | 34.2 |
| Instability/Dislocation | 3 | 8.3 | 15.8 | 181 | 2.1 | 27.1 |
| Loosening | 5 | 13.9 | 26.3 | 103 | 1.2 | 15.4 |
| Infection | | | | 44 | 0.5 | 6.6 |
| Fracture | 1 | 2.8 | 5.3 | 19 | 0.2 | 2.8 |
| Pain | | | | 15 | 0.2 | 2.2 |
| Arthrofibrosis | 1 | 2.8 | 5.3 | 12 | 0.1 | 1.8 |
| Malposition | | | | 9 | 0.1 | 1.3 |
| Wear Glenoid Insert | 2 | 5.6 | 10.5 | 9 | 0.1 | 1.3 |
| Incorrect Sizing | | | | 8 | 0.1 | 1.2 |
| Implant Breakage Glenoid | | | | 7 | 0.1 | 1.0 |
| Dissociation | | | | 6 | 0.1 | 0.9 |
| Implant Breakage Glenoid Insert | 1 | 2.8 | 5.3 | 6 | 0.1 | 0.9 |
| Lysis | 1 | 2.8 | 5.3 | 6 | 0.1 | 0.9 |
| Metal Related Pathology | | | | 6 | 0.1 | 0.9 |
| Progression Of Disease | | | | 2 | 0.0 | 0.3 |
| Other | | | | 7 | 0.1 | 1.0 |
| N Revision | 19 | 52.8 | 100.0 | 669 | 7.6 | 100.0 |
| N Primary | 36 | | | 8757 | | |


Note: This table is restricted to revisions within 10.6 years for all groups to allow a time-matched comparison of revisions. Note: Prostheses no longer used in 2021 are excluded from the comparator.

FIGURE 2

Cumulative Incidence Revision Diagnosis of Primary Total Stemmed Shoulder Replacement

This figure details the cumulative incidence of the most common reasons for revision. The five most common reasons for revision are included as long as each of these reasons account for more than 10 procedures or at least 5% of all revisions for the Vaios/Vaios total stemmed shoulder combination. A comparative graph is provided of the cumulative incidence for the same reasons for revisions for all other total stemmed shoulder prostheses.

Figure 2: Cumulative Incidence Revision Diagnosis for Primary Total Stemmed Shoulder Replacement



Type of Revision Performed for Primary Total Stemmed Shoulder Replacement

This analysis identifies the components used in the revision of the Vaios/Vaios total stemmed shoulder combination and compares it to the components used in the revision of all other total stemmed shoulder prostheses.

The reason this analysis is undertaken is to identify whether there is one or more components which are being replaced that differ from the components replaced for revisions of all other total stemmed shoulder prostheses i.e. is there a difference in the type of revision undertaken for the Vaios/Vaios total stemmed shoulder combination compared to all other total stemmed shoulder prostheses.

Table 5: Primary Total Stemmed Shoulder Replacement - Type of Revision (Follow-up Limited to 10.6 Years)

| | Vaios/Vaios | | Other Total Stemmed Shoulder | |
|---------------------------|-------------|---------|------------------------------|---------|
| Type of Revision | Number | Percent | Number | Percent |
| Humeral Component | 11 | 57.9 | 363 | 54.3 |
| Humeral/Glenoid | 3 | 15.8 | 186 | 27.8 |
| Glenoid Component | 1 | 5.3 | 28 | 4.2 |
| Cement Spacer | | | 22 | 3.3 |
| Removal of Prostheses | | | 6 | 0.9 |
| Reinsertion of Components | | | 1 | 0.1 |
| N Major | 15 | 78.9 | 606 | 90.6 |
| Head Only | 3 | 15.8 | 52 | 7.8 |
| Reoperation | | | 5 | 0.7 |
| Minor Components | 1 | 5.3 | 3 | 0.4 |
| Head/Insert | | | 2 | 0.3 |
| Cement Only | | | 1 | 0.1 |
| N Minor | 4 | 21.1 | 63 | 9.4 |
| TOTAL | 19 | 100.0 | 669 | 100.0 |

Note: This table is restricted to revisions within 10.6 years for all groups to allow a time-matched comparison of revisions. Note: Prostheses no longer used in 2021 are excluded from the comparator.

Revision Rates of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Fixation

This analysis is provided as some prostheses have more than one fixation option. Additionally there are prostheses where an alternative to the recommended approach to fixation was used e.g. a cementless prosthesis that has been cemented or vice-versa.

Table 6: Revised Number of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Fixation

| Fixation | N Revised | N Total |
|-----------------------------|-----------|---------|
| Cemented | 1 | 1 |
| Cementless | 15 | 25 |
| Hybrid (Glenoid Cemented) | 2 | 8 |
| Hybrid (Glenoid Cementless) | 1 | 2 |
| TOTAL | 19 | 36 |

Revision Rates of Primary Total Stemmed Shoulder Replacement by State

This enables a state by state variation to be identified for the Vaios/Vaios total stemmed shoulder combination and provides the comparative data for each of the states for all other total stemmed shoulder prostheses.

The purpose of this analysis is to determine if the higher than anticipated rate of revision has widespread distribution between states. If there is widespread distribution then the reason for the higher than anticipated rate of revision is unlikely to be surgeon specific. If the prosthesis has been used in only a small number of states it is not possible to distinguish if the higher than anticipated rate of revision is related to the prosthesis, surgeon, technique or patient.

Table 7: Revised Number of Primary Total Stemmed Shoulder Replacement by State

| Component | State | N Revised | N Total |
|------------------------------|--------|-----------|---------|
| Vaios/Vaios | NSW | 9 | 12 |
| | VIC | 3 | 5 |
| | QLD | 3 | 8 |
| | WA | 4 | 8 |
| | TAS | 0 | 1 |
| | ACT/NT | 0 | 2 |
| Other Total Stemmed Shoulder | NSW | 191 | 2613 |
| | VIC | 157 | 2055 |
| | QLD | 99 | 1467 |
| | WA | 134 | 1184 |
| | SA | 67 | 1015 |
| | TAS | 26 | 201 |
| | ACT/NT | 24 | 222 |
| TOTAL | | 717 | 8793 |

Number of Revisions of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Year of Implant

This analysis details the number of prostheses reported each year to the Registry for the Vaios/Vaios total stemmed shoulder combination. It also provides the subsequent number of revisions of the primaries reported in that year.

Primary procedures performed in later years have had less follow up time therefore the number revised is expected to be less than the number revised in earlier years. For example, a primary procedure performed in 2021 has a maximum of one year to be revised, whereas a primary procedure performed in 2019 has a maximum of three years to be revised.

Table 8: Number of Revisions of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Year of Implant

| Year of Implant | Number Revised | Total Number |
|-----------------|----------------|--------------|
| 2011 | 8 | 16 |
| 2012 | 9 | 17 |
| 2013 | 2 | 2 |
| 2014 | 0 | 1 |
| TOTAL | 19 | 36 |

Revision Rates of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Catalogue Number Range

Many prostheses have a number of catalogue ranges. The catalogue range is specific to particular design features; more than one catalogue range usually indicates a minor difference in design in a particular Vaios/Vaios prosthesis.

This analysis has been undertaken to determine if the revision rate varies according to the catalogue number range.

| Model | Catalogue Range | Catalogue Description | Cemen | t Material |
|--------------|-----------------|---------------------------|-------|-------------------------------|
| Humeral Stem | | | | |
| Vaios | 8010108-8010116 | STEM H-AC TI6AI4V | NO | METAL |
| Vaios | 8020106-8020112 | CEMENTED STEM | YES | METAL |
| Vaios | 8022208-8022210 | LONG DISTAL STEM CEMENTED | YES | METAL |
| Glenoid | | | | |
| Vaios | 8010900-8010911 | METAL BACK GLENOID | NO | METAL |
| Vaios | 8020342-8020458 | UHMWPE CEMENTED GLENOID | YES | NON CROSS-LINKED POLYETHYLENE |

Table 9: Revised Number of Vaios/Vaios Primary Total Stemmed Shoulder Replacement by Catalogue Number Range

| Humeral Stem Range | Glenoid Range | N Revised | N Total | |
|-----------------------|-----------------|-----------|---------|--|
| 8010108-8010116 | 8010900-8010911 | 15 | 25 | |
| | 8020342-8020458 | 3 | 9 | |
| 8020106-8020112 | 8010900-8010911 | 0 | 1 | |
| 8022208-8022210 | 8010900-8010911 | 1 | 1 | |
| TOTAL | | 19 | 36 | |